

On the origin of very high energy γ -rays from radio galaxies

Frank M. Rieger

Max-Planck-Institut für Kernphysik, P.O. Box 103980, 69029 Heidelberg, Germany

Abstract. Radio galaxies have emerged as a new gamma-ray emitting source class on the extragalactic sky. With their jets misaligned, i.e. not directly pointing towards us, they offer a unique tool to probe some of the fundamental (and otherwise hidden) non-thermal processes in AGN. This contribution briefly summarizes the observed characteristics of the four radio galaxies detected so far at very high energies (VHE). Given its prominence, particular attention is given to the origin of the *variable* VHE emission in M87. We discuss some of the theoretical progress achieved for this source within recent years highlighting, amongst others, the relevance of magnetospheric particle acceleration and emission models.

Keywords: Radio galaxies, gamma-rays, jets, black holes, physical processes

PACS: 98.54.Gr, 95.30.Qd, 95.85.Pw, 98.62.En, 98.62.Nx

INTRODUCTION

The extragalactic sky as observed at very high energies (VHE; > 100 GeV) is highly dominated by Active Galactic Nuclei (AGNs) of the blazar-type, i.e., AGNs with relativistic jets pointing almost directly towards us. Flux enhancement of the intrinsic non-thermal jet emission by relativistic Doppler boosting effects then naturally favours their detection at VHE. On the other hand, misaligned (i.e., non-blazar) AGNs, characterized by jets substantially inclined with respect to the observer, could allow us to probe some of the fundamental non-thermal VHE processes in AGN that are otherwise swamped by boosted jet emission. Nearby radio galaxies (RGs) are especially attractive as their proximity often makes it possible to resolve their radio jets down to sub-parsec scales, and to investigate the behavior of the emission at different wavelengths.

RADIO GALAXIES AT GAMMA-RAY ENERGIES

At high gamma-ray energies (HE; > 100 MeV) Fermi-LAT has reported the detection of about ten misaligned RGs, mostly of the Fanaroff-Riley-type I (FR I) [1, 2, 3], see also Fig. 1. At very high energies (VHE), only four RGs have been clearly identified so far: Cen A ($d \simeq 3.8$ Mpc), M87 ($\simeq 16.7$ Mpc) and the Perseus Cluster ($d \sim 77$ Mpc, $z \sim 0.018$) sources NGC 1275 and IC 310. A detection of the RG 3C66B at VHE was reported by MAGIC (2007 observations [4]), but the VHE emission seems not sufficiently disentangled from the nearby (separation $\theta \sim 0.12^\circ$) IBL blazar 3C66A to consider it here as a confirmed VHE emitter.

Name	Type	Distance	LAT/HE detection	VHE
Cen A	FR I	3.7 Mpc	LAT 2010	✓
M87	FR I	16 Mpc	LAT 2009	✓
Fornax A	FR I	18 Mpc	LAT 2011	
Cen B	FR I	56 Mpc	LAT 2011	
NGC 1275	FR I	75 Mpc	LAT 2009	✓
IC 310	?	80 Mpc	LAT 2010	✓
NGC 6251	FR I	106 Mpc	LAT 2010	
3C 78	FR I	124 Mpc	LAT 2010*	
3C 120	FR I	142 Mpc	LAT 2010*	
3C 111	FR 2	213 Mpc	LAT 2010*	
PKS 0943-76	FR 2	1360 Mpc	LAT 2010	
.....				

* only in 1st LAT AGN catalog (missing in 2LAC/2FGL)

FIGURE 1. Radio Galaxies (RGs) detected at gamma-ray energies. Out of ≥ 886 AGN ("clean" sample) detected by Fermi at high energies (HE) less than a few percent are RGs. Only four of these RGs are currently confirmed to be VHE emitters.

The nearest AGN *Cen A*, hosting a black hole of mass $M_{\text{BH}} \simeq (0.5 - 1) \times 10^8 M_{\odot}$, was the second RG to be detected at VHE energies in a deep ($> 120\text{h}$) exposure by H.E.S.S. [5]. *Cen A* is a weak VHE source with an integral flux $I(> 250 \text{ GeV})$ above 250 GeV of about 0.8% of the Crab Nebula (equivalent to an apparent isotropic VHE luminosity of $L(> 250 \text{ GeV}) \simeq 2 \times 10^{39} \text{ erg/s}$). The VHE spectrum extends up to $\sim 5 \text{ TeV}$ and is compatible with a single power law of photon index 2.7 ± 0.5 . No significant VHE variability has been found so far, so that an extended origin of the VHE emission [6] cannot be simply excluded yet. At high energies, Fermi-LAT has also detected HE γ -rays up to 10 GeV from the core of *Cen A* [7]. The HE light curve (15d bins) of the core is consistent with no variability and the HE γ -ray spectrum can be described by a photon index comparable to the one in the VHE. A simple power law extrapolation of the HE spectrum, however, cannot account for the observed TeV flux. This could be indicative of an additional emission contribution to the VHE domain beyond the common synchrotron-Compton emission [8], emerging at the highest energies [9]. At GeV energies, Fermi-LAT has also detected γ -ray emission from the giant radio lobes of *Cen A*. A recent analysis of a larger HE data set revealed significant evidence for a spatial extension of the γ -ray lobes beyond the radio (WMAP) image [10], suggesting

the need for a more complex ($\geq$ two-zone) lobe emission modeling than done so far. Note that the angular resolution of H.E.S.S. ($\sim 0.1^\circ$) clearly allows to exclude the lobe regions as source of the detected TeV emission.

The Virgo Cluster galaxy *M87* was the first radio galaxy detected at TeV energies [11]. Classified as an FR I, M87 is known to host one of most massive black holes with $M_{\text{BH}} \simeq (2 - 6) \times 10^9 M_\odot$ and to exhibit a relativistic jet misaligned by $\theta \simeq (15 - 25)^\circ$. These latter values are consistent with modest Doppler boosting $D = 1/[\Gamma_j(1 - \beta \cos \theta)] \lesssim 3$ (for review, see e.g. [12]). M87 has shown particularly interesting VHE features within recent years, including rapid day-scale variability (observed flux doubling time scales $\Delta t_{\text{obs}} \sim 1$) above 350 GeV during active source states (with flux levels sometimes exceeding 10% of the Crab Nebula), and a hard spectrum (compatible with a power law of photons index $\Gamma \simeq 2.2$) extending beyond 10 TeV [13, 14, 15, 16, 17]. Both the hard VHE spectrum and the observed rapid VHE variability are remarkable features for a misaligned AGN. M87 is the only RG where clear evidence for day-scale TeV variability (see Fig. 2) has been found, although there appear to be some hints for day-scale variability in IC 310 (and for month-type in NGC 1275) as well (see below). The observed variations in M87 seen at VHE are the fastest compared to any other waveband so far, and imply a compact γ -ray emitting region ($R < c\Delta t_{\text{obs}}D$), comparable to the Schwarzschild radius $r_s = (0.6 - 1.8) \times 10^{15}$ cm of its black hole. This could point to the black hole vicinity as the most likely origin of the observed, variable VHE radiation, a consideration supported by high-resolution radio (VLBA) observations during the 2008 (February) VHE high state of M87, showing that the VHE outburst was followed by an ejection of a new radio component from very close to the black hole [18].

The Perseus Cluster galaxy *IC 310* has been detected by MAGIC above 300 GeV in about 21h of data (taken in 2009/2010) at an average level of $\sim 3\%$ of the Crab Nebula [19]. The source has been originally classified in the literature as a head-tail RG, but a recent radio analysis finds little evidence for jet bending [20] (see also Ref. [21] for the proposal that IC 310 may instead be a weakly beamed blazar). The VHE spectrum measured between 150 GeV and 7 TeV is very hard (even harder than in M87) and compatible with a single power law of photon index $\Gamma \simeq 2.0$. There is clear evidence for VHE variability on yearly and monthly time scales, with some indications for day-scale activity found in a new analysis [22], features that are reminiscent of the VHE activity seen in M87. Note however, that due to its larger distance, IC 310 needs to be intrinsically much more luminous.

In addition to IC 310, MAGIC has also recently reported the detection of the central dominant (FR I) cluster galaxy *NGC 1275* above ~ 100 GeV during enhanced high energy (Fermi-LAT) activity in 46h of data (taken between 08/2010-02/2011) [23]. This source is known to have radio jets misaligned by $\gtrsim 30^\circ$. Fermi-LAT data indicate significant flaring activity above 0.8 GeV down to time scales of days [25]. At VHE energies the situation is less evident, with no evidence for VHE variability found in the noted data set, while a recent, improved analysis of an earlier (10/2009-02/2010) VHE data set provides hints for a possible month-type variability [24]. NGC1275 shows a very steep VHE spectrum (power law photon index $\Gamma \simeq 4.1$) extending up to ~ 500 GeV. When compared with the hard HE (Fermi-LAT) spectrum (photon index $\Gamma \simeq 2.1$), this implies the existence of a break or cut-off in the SED around some tens of GeV.




FIGURE 2. The 2010 April VHE flare of the radio galaxy M87 as observed by different Cherenkov Telescopes. Significant day-scale activity is evident. The curve shows a fit of an exponential function to the data. From Ref. [16].

ON THE ORIGIN OF VARIABLE TEV GAMMA-RAYS FROM M87

Being one of the best studied extragalactic object, M87 has become a prominent benchmark for theoretical research. A multitude of VHE emission scenarios, operating on different scales and relying on either hadronic and/or leptonic radiation processes, have been developed and applied within recent years (cf. Fig. 3). The observed day-scale variability favors an origin of the variable TeV emission on scales of the inner jet and below, although outer-jet scenario may possibly contribute to the overall quiescent VHE source state (e.g., [12] for discussion).

(1) One recent theoretical development in the context of leptonic models considers relativistic (Petschek-type) reconnection as acceleration mechanism of electrons to high energy [26], see Fig. 4 for illustration. Compton up-scattering of external photons could then possibly account for the production of the observed, variable VHE gamma-rays [27]. Efficient reconnection in a highly magnetized ($\sigma \sim 100$) electron-proton jet could lead to an additional relativistic velocity component of the ejected plasma ($\Gamma_r \simeq \sqrt{\sigma}$, at some angle to the main outflow direction) with respect to the mean bulk flow of the jet. This makes strong, differential Doppler boosting effects for the emitting zone possible, circumventing the problem of a modest Doppler factor for the general bulk flow. At present, there seem to be a number of open issues that need to be clarified within such an approach in order to assess its explanatory potential. This includes the fact that (i) typically much lower magnetizations ($\sigma \sim 10$) are expected for electron-proton (disk-driven) jets in AGN (in contrast to BZ-driven electron-positron jets), that (ii) at the anticipated

HST-I	EC starlight photons (e.g. Stawarz+06)	
inner jet (sub-parsec)	leptonic	decelerating flow (e.g. Georgopoulos+05)
		spine-shear (e.g. Tavecchio+08)
		mini/multi-blobs (e.g. Lenain+08)
		reconnection (e.g. Giannios+10)
	hadronic	proton synchrotron & p- γ (e.g. Reimer+04)
		jet-star interactions / pp (e.g. Barkov+12)
		combined lepto-hadronic (e.g. Reynoso+11)
Magnetosphere	rotational acceleration & IC (e.g. Rieger & Aharonian 08)	
	gap-type particle acceleration & IC (e.g. Levinson & Rieger 11)	

FIGURE 3. Possible theoretical scenarios for the origin of the variable VHE emission as proposed for M87. For details, including some pros and cons of the models, see Ref. [12].

scale of VHE production a non-negligible magnetic guide field may persist allowing for weak dissipation only, and that (iii) the achievability of power-law electron acceleration much beyond the thermal Lorentz factor $\sqrt{\sigma} m_p / 2m_e$ is not clear.

(2) In the context of hadronic models, on the other hand, one new development considers inelastic proton-proton (pp) collisions (and π^0 -decay) in the jet for the origin of the observed VHE emission. While AGN jets have been often considered not to carry (on average) a sufficient amount (n_p) of target matter [28, 29] to ensure the efficiency of this (pp) process (given its characteristic timescale $t'_{pp} \simeq 10^{15}/n'_p$ sec), it seems possible that interactions of a red giant star or a massive, dense gas cloud (size r_c) with the base of the jet (radius r_j) could occasionally introduce the required amount of matter and thereby potentially drive fast VHE activity [30, 31]. Recent calculations suggest that such a scenario could account for the observed temporal and spectral properties if the jet is powerful enough and a sufficiently large fraction ($\propto r_c^2/r_j^2$) of it can be channeled into VHE γ -ray production. Given the observed large opening angle and transversal dimension of the radio jet in M87, models of this type require a spine-shear-type jet configuration with most of the energy flux concentrated into a narrow core.

(3) An alternative explanation has been pursued within magnetospheric models (see e.g. [32] for review), motivated by the observed day-scale VHE variability and the observed




FIGURE 4. LEFT: Illustration of a jets-in-jet scenario [26], where a variety of ‘mini-jet’ features are thought to be induced by relativistic magnetic reconnection events within the general jet flow. This could lead to an additional velocity component (Γ_r) relative to the main flow and allow a favorable orientation with respect to the observer. RIGHT: Illustration of a hadronic model, where interactions of the jet with a massive gas cloud leads to Fermi (shock)-type proton acceleration and introduces a sufficient target density to allow for efficient pp -collisions [31].

radio-VHE connection in 2008. For a black hole mass in M87 expected to be in the range $M_{\text{BH}} \simeq (2 - 6) \times 10^9 M_{\odot}$ [12], anticipated variability timescales could possibly be as small as $\sim r_g/c = 0.1$ d, allowing to accommodate the observed rapid VHE variability. One of the latest developments in this context analyses the possible link between VHE γ -ray activity and jet formation [33]. A schematic representation is shown in Fig. 5:

In this scenario, a maximally rotating black hole, which is embedded in an external magnetic field B generates an electric potential difference across a (vacuum) gap of height h , given by $\Delta V_e = 1.7 \times 10^{20} B_3 M_9 (h/r_s)^2$ Volts, where $B_3 = B/10^3$ G and $M_9 = M_{\text{BH}}/10^9 M_{\odot}$. Pair creation in a hot accretion flow (i.e. by $\gamma\gamma$ -annihilation of MeV photons in an ADAF or RIAF) then leads to the injection of primary electrons and positrons into the gap, with the charged particles being quickly accelerated to high Lorentz factors. The number density of ADAF bremsstrahlung MeV photons is $n_{\gamma} \propto \dot{m}^2$, so that the e^+e^- -injection rate inside the magnetosphere roughly becomes $n_{\pm} \sim \sigma_{\gamma\gamma} n_{\gamma}^2 r_s \propto \dot{m}^4$. When expressed in terms of the Goldreich-Julian (GJ) density $n_{\text{GJ}} = \Omega B / (2\pi e c) \propto \dot{m}^{1/2}$, the ratio becomes [33]

$$n_{\pm}/n_{\text{GJ}} \simeq 0.06 \dot{m}_{-4}^{7/2} M_9^{1/2}.$$

This numerical estimate has some uncertainty as the bremsstrahlung spectrum is dependent on T_e , which itself depends on the accretion rate $\dot{m} \sim 10^{-4}$ (expressed in terms of the Eddington rate). The sensitive dependence on $\dot{m}$ suggests, however, that a gap (with $n_{\pm}/n_{\text{GJ}} < 1$) can be formed during periods of low accretion. Gap-type particle acceleration and subsequent curvature emission and, in particular, inverse Compton (IC) up-scattering of disk photons then produces γ -rays with a spectrum extending up to 10^4 TeV. Different contributions dominating at different energy bands could then lead to a different variability behavior as e.g. seen by MAGIC during the 02/2008-VHE high




FIGURE 5. VHE γ -ray production within a black hole-magnetospheric scenario [33]: Electrons and positrons are accelerated in a vacuum gap to high Lorentz factors. The gap is exposed to soft radiation field of the disk. Curvature emission and inverse Compton up-scattering of ambient disk photons yields VHE γ -rays and triggers e^+e^- -cascades. The resultant large pair multiplicity establishes the force-free outflow seen in VLBA images.

state in M87 [14]. For low accretion rate, VHE photons with energies below several TeV can escape $\gamma\gamma$ -absorption (see e.g., [33, 12] for details). On the other hand, IC photons with energies above ~ 10 TeV interact with the ambient radiation field initiating pair cascades just above the gap. This leads to a large pair multiplicity ($M = n_{2\pm}/n_{\text{GJ}}$ up to 10^3). A force-free outflow becomes established above the pair formation front and accounts for the jet feature seen in radio VLBA. Given the sensitive dependence of gap formation (primary pair injection) and photon escape (optical depth for $\gamma\gamma$ -absorption) on accretion rate, any emission from the gap is likely to be intermittent. Modest changes in accretion rate, for example, may naturally give rise to the variability of the TeV emission and to fluctuations of the resultant force-free outflow.

CONCLUSIONS

Radio galaxies have emerged as a new and particularly interesting VHE source class on the extragalactic sky. Evidence for γ -ray variability, as seen in most of the sources (apart from Cen A) suggests that the observed γ -ray emission originates in a compact region. In reality, the situation may be somewhat more complex with extended regions contributing to the average/low states of the source, and different (additional) zones dominating during high VHE source states. In the case of M87, the observed VHE flaring characteristics (such as, e.g., the hard γ -ray spectrum extending beyond 10 TeV and the day-scale VHE activity) are of fundamental interest as they allow us to probe the near

black hole environment and to study the possible link between VHE activity and jet formation in AGN. Radio galaxies like M87 thus appear to be particularly promising targets for future & more sensitive instruments like CTA.

ACKNOWLEDGMENTS

Discussions with Felix Aharonian, Amir Levinson, Maxim Barkov, Matthias Beilicke and Martin Raue, and financial support by a LEA Fellowship are gratefully acknowledged.

REFERENCES

1. A. Abdo et al. (Fermi-LAT Collaboration), *ApJ* 720, 912 (2010)
2. M. Ackermann et al. (Fermi-LAT Collaboration), *ApJ* 743, 171 (2011)
3. P.L. Nolan et al. (Fermi-LAT Collaboration), *ApJS* 199, 31 (2012)
4. E. Aliu et al. (MAGIC Collaboration), *ApJL* 692, L29 (2009)
5. F. Aharonian et al. (H.E.S.S. Collaboration), *ApJL* 695, L40 (2009)
6. M. Hardcastle, these proceedings (2012)
7. A.A. Abdo et al. (Fermi-LAT Collaboration) *ApJ* 719, 1433 (2010)
8. M. Chiaberge et al., *MNRAS* 324, 33 (2001)
9. F.M. Rieger, *Mem. S.A.It.* 83, 127 (2012)
10. R-Z. Yang et al., *A&A* 542, A19 (2012)
11. F. Aharonian et al. (HEGRA Collaboration), *A&A* 403, L1 (2003)
12. F.M. Rieger and F. Aharonian, *MPLA* 27, 12300301 (2012)
13. F. Aharonian et al. (H.E.S.S. Collaboration), *Science* 314, 1424 (2006)
14. J. Albert et al. (MAGIC Collaboration), *ApJ* 685, L23 (2008)
15. E. Aliu et al. (VERITAS Collaboration), *ApJ* 746, 141 (2012)
16. A. Abramowski et al., *ApJ* 746, 151 (2012)
17. M. Beilicke et al., these proceedings
18. V.A. Acciari et al. (VERITAS, MAGIC, VLBA M87 and H.E.S.S. Collab.), *Science* 325, 444 (2009)
19. J. Aleksic et al. (MAGIC Collaboration), *ApJL* 723, 207 (2010)
20. M. Kadler et al., *A&A* 538, L1 (2012)
21. T.A. Rector, J.T. Stocke and E.S. Perlman, *ApJ* 516, 145 (1999)
22. D. Eisenacher, P. Colin et al., these proceedings (2012)
23. J. Aleksic et al. (MAGIC Collaboration), *A&A* 539, 2 (2012)
24. P. Colin, D. Eisenacher et al., these proceedings (2012)
25. A.M. Brown and J. Adams, *MNRAS* 413, 2785 (2011)
26. D. Giannios, D.A. Uzdensky and M.C. Begelman, *MNRAS* 402, 1649 (2010).
27. Y-D. Cui et al., *ApJ* 746, 177 (2012).
28. A. Celotti et al., *MNRAS* 293, 288 (1998).
29. M. Wardle et al., *Nature* 395, 457 (1998).
30. M.V. Barkov, F.A. Aharonian and V. Bosch-Ramon, *ApJ* 724, 1517 (2010).
31. M.V. Barkov, V. Bosch-Ramon and F.A. Aharonian, *ApJ* 755, 170 (2012).
32. F.M. Rieger, *IJMPD* 20, 1547 (2011).
33. A. Levinson and F. Rieger, *ApJ* 730, 123 (2011).